

plantmate™

drench

TRICHO PROTECTION™
healthy plants

Rootzone booster WP

Recommended for all growing plants

**Beneficial
Bio-inoculant**

A WP formulation for application to the root zones of all plants via all systems including conventional dripper, tape, overhead irrigation and boom spray.

Active ingredient:

Proprietary strains of *Trichoderma harzianum* 100 million cfu/g — selected natural strains formulated in the specific colonisation matrix.



plantmate™

1.5kg net

plantmate™ drench

Instructions for use

Ensure the mixing tank has been thoroughly washed to remove any chemical residues before mixing plantmate drench with fresh clean water. Always pre-mix plantmate drench by adding the powder to water at the minimum rate of 500g to 5 litres. Mix well and allow to stand for 5 minutes before adding to the dispensing tank, which should contain a minimum of 100 litres. Ensure the dispensing tank is agitated continuously during application. After applications through micro delivery systems, dripper & tape etc., always flush with clean water. Plantmate drench may be applied with fertilizers, seaweed extracts and fish-based products.

Indoor and Glasshouse crops

Organic media (sawdust, peat, coir, bark based growing media). Apply plantmate drench at the minimum rate of 500g per 5000 square metres. Apply every 10–14 days.

Inorganic media (rockwool, pumice, vermiculite & other artificial substrates). Apply plantmate drench at the minimum rate of 500g per 1000 litres. Apply every 10–14 days.

Hydroponic/NFT/Aeroponic Apply plantmate drench at the minimum rate of 500g per 1000 litres. Apply every 10–14 days. These recommendations apply to all systems including, open, closed, re-circulating and run to waste.



Manufactured under licence for Agrimm Proprietary Ltd,
58 Coolamon Close, Ocean Grove, VIC 3226, Australia
Tel: 1800 42 47 46 Fax: 03 5255 4506
info@agrimm.com www.agrimm.com

Manufactured by Agrimm Technologies Ltd,
Christchurch, New Zealand.

Field crops, potatoes

Apply plantmate drench at the minimum rate of 750g per hectare through 'T' tape, drip, or other irrigation systems. Apply every 4 weeks or more often if the crop is under stress from difficult growing conditions. For potatoes spray plantmate drench in furrow at planting.

Nurseries, seed planting, seedling plugs

Apply plantmate drench at the minimum rate of 250g per 100 litres onto the seed bed. Apply every 4 weeks. For seedling plugs, prior to release for field planting, apply 500g per 100 litres.


Orchards and vineyards

Apply plantmate drench at the rate of 500g-750g per hectare. Apply 3 to 5 times per season depending upon climate, crop stress and growing conditions. Apply with nutrient support as recommended in the treatment programme. Application is recommended during plant establishment, prior to periods of new growth flush and whilst carrying heavy crop loading.

Precautions for use

- Keep out of reach of children
- Store refrigerated at 4°C
- To avoid airborne particles wear mask when dispensing product
- Wash hands and exposed skin after use and before meals
- Crush the empty sachet and bury in approved landfill.
- Do not contaminate water supply with material or the empty pack.

NOTICE TO BUYER: The manufacturer of plantmate drench makes no warranty of any kind on the use of this product and accepts no liability for results from its use whether used in accordance with the label directions or otherwise.



9 421900 077021

www.agrimm.co.nz



BEST BEFORE: 1 2 3 4 5 6 7 8 9 10 11 12 /16 17 18 19 20

Batch :